

After Your Metacarpal Fracture Surgery

Dr. Paul Nielsen

Dressings and Wound Care

- After surgery you will have a bulky splint or wrap on your hand to protect it
- Leave the splint or dressing in place until follow up
 - You must cover the splint for showering to keep it dry
- Call our office at 308-865-2570 if something happens to your splint or wrap

Pain Medicine

- NSAIDs such as ibuprofen or naproxen are recommended to improve pain after surgery in addition to prescription pain medication
- You will be prescribed a strong pain medication, such as hydrocodone and oxycodone, to control your pain after surgery
 - You should start taking pain medication as soon as you start getting feeling back, indicating the nerve block is wearing off to avoid rebound pain
 - You are allowed to drive when you are not taking prescription pain medication

Activity

- You are encouraged to move the exposed part of your fingers immediately after surgery
 - The knuckle closest to the fracture may be held still in the splint
- Elevate your hand above your elbow and heart for the first few days after surgery
- Do not lift anything with your arm until after your first post-operative appointment

Follow-Up Care

- You will have an appointment in clinic 5-7 days after surgery to switch to a removable splint
 - Additional appointments vary depending on the exact procedure
- Hand therapy is sometimes required if stiffness develops from 2 to 6 weeks after surgery

Results

- Soreness and stiffness often persist for a few months after surgery
- Most patients regain a functional range of motion after surgery
- Hardware used to fix the break occasionally must be removed to improve range of motion